

One-loop calculation of mass dependent $\mathcal{O}(a)$ improvement coefficients for the relativistic heavy quarks on the lattice

Sinya Aoki^a, Yasuhisa Kayaba^a and Yoshinobu Kuramashi^b

^aInstitute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan

^bHigh Energy Accelerator Research Organization(KEK), Tsukuba, Ibaraki 305-0801, Japan

We carry out the one-loop calculation of mass dependent $\mathcal{O}(a)$ improvement coefficients in the relativistic heavy quark action recently proposed, employing the ordinary perturbation theory with the fictitious gluon mass as an infrared regulator. We also determine renormalization factors and improvement coefficients for the axial-vector current at the one-loop level. It is shown that the improvement coefficients are infrared finite at the one-loop level if and only if the improvement coefficients in the action are properly tuned at the tree level.

1. INTRODUCTION

While, in principle, lattice QCD allows a precise determination of weak matrix elements associated with the B and D mesons, c - and b -quarks are too heavy to treat directly in current numerical simulations due to large $\mathcal{O}(m_Q a)$ errors.

Recently, a new relativistic approach to control $m_Q a$ errors was proposed from the view point of the on-shell $\mathcal{O}(a)$ improvement program assuming $m_Q \gg \Lambda_{\text{QCD}}$ [1,2]. The action is given by

$$\begin{aligned} S_q = & a^4 \sum_x \bar{q}(x) [\gamma_0 D_0 + \nu \gamma_i D_i + m_0 \\ & - ar_t/2D_0^2 - ar_s/2D_i^2 - aigc_B/4\sigma_{ij}F_{ij} \\ & - aigc_E/2\sigma_{0i}F_{0i}] q(x), \end{aligned} \quad (1)$$

where we are allowed to choose $r_t = 1$ and other four parameters ν , r_s , c_E and c_B are analytic functions of $m_Q a$ and the gauge coupling g . Once a nonperturbative adjustment of four parameters ν , r_s , c_E and c_B is achieved, the remaining cutoff effects are reduced to be $\mathcal{O}((a\Lambda_{\text{QCD}})^2)$. In this report we present a perturbative determination of c_E and c_B up to the one-loop level (see Ref. [2] for details). In addition we calculate the renormalization factors and the $\mathcal{O}(a)$ improvement coefficients for the axial-vector current at the one-loop level.

2. DETERMINATION OF c_E AND c_B AT THE ONE-LOOP LEVEL

We determine $c_E = c_E^{(0)} + g^2 c_E^{(1)}$ and $c_B = c_B^{(0)} + g^2 c_B^{(1)}$ from the on-shell quark-quark scattering amplitude. In the massless case, c_{SW} is successfully determined up to the one-loop level employing the conventional perturbation theory with the fictitious gluon mass as an infrared regulator [3]. We extend this calculation to the massive case. Since the tree level determination of c_E and c_B in the massive case is already performed in Ref. [1], next step is to calculate six types of one-loop diagrams depicted in Fig. 1.

We first consider to calculate $c_B^{(1)}$. Without the space-time symmetry the general form of the off-shell vertex function at the one-loop level is written as

$$\begin{aligned} \Lambda_k^{(1)}(p, q, m) = & \gamma_k F_1^k + \gamma_k \{\not{p} F_2^k + \not{p}_0 F_3^k\} + \{\not{q} F_4^k + \not{q}_0 F_5^k\} \gamma_k \\ & + \not{q} \gamma_k \not{p} F_6^k + \not{q} \gamma_k \not{p}_0 F_7^k + \not{q}_0 \gamma_k \not{p} F_8^k \\ & + (p_k + q_k) [H_1^k + \not{p} H_2^k + \not{q} H_3^k + \not{q} \not{p} H_4^k] \\ & + (p_k - q_k) [G_1^k + \not{p} G_2^k + \not{q} G_3^k + \not{q} \not{p} G_4^k] + \mathcal{O}(a^2), \end{aligned} \quad (2)$$

where $\Lambda_k = \Lambda_k^{(0)} + g^2 \Lambda_k^{(1)} + \mathcal{O}(g^4)$ and $\not{p} = \sum_{\alpha=0}^3 p_\alpha \gamma_\alpha$, $\not{q} = \sum_{\alpha=0}^3 q_\alpha \gamma_\alpha$, $\not{p}_0 = p_0 \gamma_0$, $\not{q}_0 = q_0 \gamma_0$. p and q are incoming and outgoing quark




Figure 1. One-loop quark-gluon vertex diagrams.

momenta, respectively. The coefficients F_i^k , G_i^k and H_i^k are functions of p^2 , q^2 , $p \cdot q$ and m . Sandwiching $\Lambda_k^{(1)}(p, q, m)$ by the on-shell quark states $u(p)$ and $\bar{u}(q)$, which satisfy $\not{p}u(p) = im_p u(p)$ and $\bar{u}(q)\not{q} = im_p \bar{u}(q)$, the matrix element is reduced to

$$\begin{aligned} & \bar{u}(q)\Lambda_k^{(1)}(p, q, m)u(p) \\ &= \bar{u}(q)\gamma_k u(p) \{F_1^k + im_p 2F_2^k - m_p^2 F_6^k\} \\ &+ (p_k + q_k)\bar{u}(q)u(p) \{H_1^k + im_p 2H_2^k - m_p^2 H_4^k\} \\ &+ \mathcal{O}(a^2), \end{aligned} \quad (3)$$


where we use the relations among F_i^k , G_i^k and H_i^k constrained by the charge conjugation symmetry.

The relevant term for the determination of c_B is the last one in eq.(3), whose coefficient can be extracted by setting $p = p^* \equiv (p_0 = im_p, p_i = 0)$ and $q = q^* \equiv (q_0 = im_p, q_i = 0)$ in eq.(2):

$$\begin{aligned} & [H_1^k + im_p(H_2^k + H_3^k) - m_p^2 H_4^k]_{p=p^*, q=q^*}^{\text{latt}} \\ &= \frac{1}{8} \text{Tr} \left[\left\{ \frac{\partial}{\partial p_k} + \frac{\partial}{\partial q_k} \right\} \Lambda_k^{(1)}(p^*, q^*, m)(\gamma_4 + 1) \right. \\ & \left. - \left\{ \frac{\partial}{\partial p_i} - \frac{\partial}{\partial q_i} \right\} \Lambda_k^{(1)}(p^*, q^*, m)(\gamma_4 + 1)\gamma_i \gamma_k \right]^{i \neq k}. \end{aligned} \quad (4)$$

We should remark that the second term in eq.(3) contains both the lattice artifact of $\mathcal{O}(p_k a, q_k a)$ and the physical contribution of $\mathcal{O}(p_k/m, q_k/m)$. The parameter c_B is determined to eliminate the lattice artifact of $\mathcal{O}(p_k a, q_k a)$:

$$\frac{c_B^{(1)} - r_s^{(1)}}{2} = [H_1^k + im_p 2H_2^k - m_p^2 H_4^k]_{p=p^*, q=q^*}^{\text{latt}}$$

Figure 2. $c_E^{(1)}$ and $c_B^{(1)}$ with the plaquette gauge action as a function of pole mass $m_p^{(0)}$.

$$-Z_q^{(0)} [H_1^k + im_p 2H_2^k - m_p^2 H_4^k]_{p=p^*, q=q^*}^{\text{cont}}.$$

The infrared behavior of eq.(4) is investigated by expanding the inner loop momentum in Fig. 1 around zero. The infrared divergence is found to be

$$\left(-\frac{1}{2N_c} + \frac{N_c}{2} \right) (c_B^{(0)} - r_s^{(0)})L - \left(-\frac{N_c}{2} \right) \frac{Z_q^{(0)}}{m_p^{(0)}}L$$

where $L = -1/(16\pi^2) \ln |\lambda^2 a^2|$ with λ the fictitious gluon mass. If the tree level improvement coefficients are properly tuned as $c_B^{(0)} = r_s^{(0)}$, we are left with $-(N_c/2)(Z_q^{(0)}/m_p^{(0)})L$, which is exactly the same as the infrared divergence in the continuum theory with the correct normalization factor $Z_q^{(0)}$.

In the same way, $c_E^{(1)}$ is determined free from the infrared divergence. Results for $c_B^{(1)}$ and $c_E^{(1)}$ with the plaquette action are plotted in Fig. 2. As expected $c_B^{(1)}$ and $c_E^{(1)}$ approaches to the massless value for $c_{\text{SW}}[4]$ as $m_p^{(0)} a$ vanishes.

3. $\mathcal{O}(a)$ IMPROVEMENT OF THE AXIAL-VECTOR CURRENT

We consider the heavy-light axial-vector current A_μ . Without space-time symmetry, the renormalized operator $A_\mu^{\text{latt}, R}(x)$ with the $\mathcal{O}(a)$ improvement is written as

$$A_\mu^{\text{latt}, R}(x) = Z_A^{\text{latt}} [\bar{q}(x)\gamma_\mu \gamma_5 Q(x)]$$




Figure 3. Δ_{A_μ} in heavy-light system with the plaquette gauge action as a function of heavy quark pole mass $m_Q^{(0)}$.

$$\begin{aligned}
 & -g^2 c_{A_\mu}^+ \partial_\mu^+ \{ \bar{q}(x) \gamma_5 Q(x) \} - g^2 c_{A_\mu}^- \partial_\mu^- \{ \bar{q}(x) \gamma_5 Q(x) \} \\
 & -g^2 c_{A_\mu}^L \{ \vec{\partial}_i \bar{q}(x) \} \gamma_i \gamma_\mu \gamma_5 Q(x) \\
 & -g^2 c_{A_\mu}^H \bar{q}(x) \gamma_\mu \gamma_5 \gamma_i \{ \vec{\partial}_i Q(x) \} \Big] \quad (5)
 \end{aligned}$$

where $Z_{A_\mu}^{\text{latt}}$ and $c_{A_\mu}^{(+,-,H,L)}$ depend on the quark masses m_Q and m_q . ∂_μ^+ and ∂_μ^- are defined as $\partial_\mu^+ = \vec{\partial}_\mu + \overleftarrow{\partial}_\mu$ and $\partial_\mu^- = \vec{\partial}_\mu - \overleftarrow{\partial}_\mu$. With the aid of equation of motion we can choose $c_{A_0}^H = c_{A_0}^L = 0$.

To extract $Z_{A_\mu}^{\text{latt}}$ and $c_{A_\mu}^{(+,-,H,L)}$ at the one-loop level from the general form of the vertex function, we have to use both decay and scattering processes. Except for the physical contribution, all parameters are infrared finite, once $\nu^{(0)}$, $r_s^{(0)}$, $c_E^{(0)}$ and $c_B^{(0)}$ are properly tuned. In Figs. 3, 4, we show numerical results for Δ_{A_μ} and $c_{A_\mu}^{(+,-,H,L)}$ in heavy-light system. The former is defined by $Z_A^{\text{latt}}/Z_A^{\text{MS}} = \sqrt{Z_Q^{(0)}}(1 - g^2 \Delta_{A_\mu})$, where we take $m_q = 0$ for the light quark. We observe anticipated features that $c_{A_\mu}^{(-,H,L)}$ vanishes as the heavy quark mass decreases, while Δ_{A_μ} and $c_{A_\mu}^+$ become close to their corresponding massless values obtained in Refs. [4,5].

This work is supported in part by the Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology. (Nos. 13135204, 14046202, 15204015, 15540251, 15740165.)




Figure 4. $c_{A_0}^{(+,-)}$, $c_{A_k}^{(+,-,H,L)}$ in heavy-light system with the plaquette gauge action

REFERENCES

1. S. Aoki, Y. Kuramashi and S. Tominaga, Prog. Theor. Phys. **109** (2003) 383.
2. S. Aoki, Y. Kayaba and Y. Kuramashi, hep-lat/0309161.
3. S. Aoki and Y. Kuramashi, hep-lat/0306015.
4. R. Wohlert, DESY preprint 87-069 (1987), unpublished; M. Lüscher and P. Weisz, Nucl. Phys. **B479** (1996) 429; S. Aoki *et al.*, Nucl. Phys. **B540** (1999) 501.
5. S. Aoki *et al.*, Phys. Rev. **D58** (1998) 074505; Y. Taniguchi and A. Ukawa, Phys. Rev. **D58** (1998) 114503.